

Radiology Corner Case #11

Coronary Artery Fistula

Guarantor: Vincent B. Ho, MD

Contributors: Cpt. Rey D. Gumboc, MC, USAR*; Maureen N. Hood, MS, RN*[†]; USN; Vincent B. Ho, MD*[†]

History

A 38-year-old man with a single episode of vague sub-sternal chest pain was referred for evaluation of possible coronary artery disease. His medical history was significant for hypertension and type-II diabetes mellitus. The patient had no prior history of thoracic trauma or surgery. During an exercise stress test, the patient reported chest pain; however, no electrocardiogram (EKG) changes were documented. A cardiac MRI axial black blood (Figure A) and black blood oblique (Figure B), and coronary x-ray angiogram oblique projections (Figures C and D) were performed.

Imaging Findings

MR imaging was performed using cardiac-gated, “black blood” double inversion recovery fast spin echo pulse sequence (DIR FSE). On DIR FSE images (axial, Figure 1A; oblique, Figure 1B), the left circumflex artery was identified by its black central lumen and noted to arise normally from the left main coronary artery. However, it was also noted to be ectatic and dilated along its course especially in its mid to distal segments within the left atrio-ventricular groove. Most importantly, the left circumflex coronary artery was noted to drain directly into the coronary sinus, consistent with the diagnosis of a left circumflex coronary artery to coronary sinus fistula. The presence and course of the fistula was confirmed on conventional x-ray angiogram (Figures 1C and 1D).

Diagnosis

Coronary Artery Fistula




Figure 1A

Figures 1A. Axial black-blood double inversion recovery fast spin echo (DIR FSE; select images from top left through bottom right representing cranial to caudal) images through the thorax demonstrates normal origin of the left main coronary artery from the aorta and normal branching of the left anterior descending coronary artery (arrow head) from the left main coronary artery. On these images, the actual origin of the left circumflex coronary artery could not be confirmed; however, the majority of the left circumflex coronary artery (arrow) could be traced from its proximal to distal segments within the left atrio-ventricular groove. In particular, the left circumflex is noted to dilated distally (bottom images) and drain via the cavernous sinus into the right atrium (RA). Ao = ascending aorta; P = pulmonary artery

**School of Medicine (AM) and Department of Radiology and Radiological Sciences (MNH, WRC,VBH), Uniformed Services University of the Health Sciences, 4301 Jones Bridge Road, Bethesda, Maryland 20814 and [†]Department of Radiology, National Naval Medical Center, 8901 Wisconsin Avenue, Bethesda, Maryland*


This is the full text version of the radiology corner case published in April 2007.

Full Text Version at: <http://rad.usuhs.mil/amsus.html>.


The opinions or assertions contained herein are the private views of the authors and are not to be construed as official or reflecting the views of the Uniformed Services University of the Health Sciences or the Department of Defense.

Reprint & Copyright © by Association of Military Surgeons of U.S., 2007.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE APR 2007		2. REPORT TYPE		3. DATES COVERED 00-00-2007 to 00-00-2007	
4. TITLE AND SUBTITLE Coronary Artery Fistula				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Uniformed Services University of the Health Sciences, Department of Radiology and Radiological Sciences, 4301 Jones Bridge Road, Bethesda, MD, 20814				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 3	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

**Figure 1B**

Figures 1B. Double-oblique "short axis" black-blood DIR FSE images in a plane parallel to the left atrioventricular groove (top left to bottom right representing base to apex). The left circumflex coronary artery (arrowhead) is seen arising from the left main coronary artery (long thinner arrow, bottom left image) and draining into the coronary sinus (short wider arrow, top left image). The proximal and distal left circumflex coronary arteries were tortuous and dilated; but the mid left circumflex coronary artery was less well seen because of volume averaging with the left atrium and pulmonary venous confluence which were also dark. Ao = ascending aorta

**Figure 1C****Figure 1D**

Figures 1C-D. Select oblique conventional x-ray views (C, early phase; D, late phase) from a coronary catheterization study following injection of iodinated contrast into the left main coronary artery. On the early phase image (C), the large and tortuous left circumflex coronary artery (arrow) is seen coursing posteriorly overlying the posterior cardiac silhouette and the left anterior descending coronary artery (arrowhead) is noted to follow its normal course anteriorly. On the late phase image (D) the dilated distal left circumflex (arrows) is noted to communicate directly with the coronary sinus (CS), which drains into the right atrium (not shown on image).

Discussion

Coronary artery fistulas are uncommon; incidentally reported in 0.1% to 0.3% of patients undergoing coronary or cardiac angiography.¹⁻³ However, of coronary artery anomalies, coronary artery fistulas are one of the most common to be hemodynamically (and clinically) significant congenital coronary artery anomalies.^{4,5} Other congenital arterial lesions that can cause abnormalities of myocardial perfusion include anomalous coronary arteries (notably, those with inter-arterial course between the ascending aorta and pulmonary artery) and congenital coronary artery stenosis.

Most coronary artery fistulas are congenital but may also occasionally result from trauma such as gunshot, shrapnel, or stab injuries; or as a complication of a surgical procedure, such as coronary bypass surgery, mitral valve replacement, or myocardial biopsies after cardiac transplantation.^{2,3,6,7}

Slightly more than half of all coronary artery fistulas involve the right coronary artery.^{2-5,8,9} In a few cases (roughly 5%^{4,5,9-11}), both left and right coronary arteries are involved. In roughly 85-90% of cases, coronary artery fistulas drain to the right side of the heart (i.e. coronary sinus, right atrium, right ventricle or pulmonary artery),^{2-5,8-11} thereby resulting in a left-to-right shunt. If the left-to-right shunt is large, blood may preferentially flow to the coronary fistula at the expense of the other coronary arteries, a phenomenon also known as "coronary steal." Coronary steal is particularly evident during times of stress and can result in myocardial ischemia to territories not supplied by the arterial fistula.^{5,8,13}

Approximately half of individuals with a coronary artery fistula are asymptomatic, typically incidentally detected

during coronary angiography.^{5, 8, 9} Though small fistulas can often go undetected in elderly patients, the likelihood and severity of symptoms generally increases with age if there is a left-to-right shunt. Symptoms when present can include angina, congestive heart failure, exertional dyspnea, bacterial endocarditis, and, even, acute myocardial infarction or spontaneous rupture with subsequent cardiac tamponade.^{5, 8, 9, 12} The typical clinical sign for a coronary fistula is often non-specific, represented by a continuous heart murmur whose qualities vary based upon its location and size.^{1, 2, 5, 8, 9, 12}

On chest radiographs, patients may have cardiomegaly or an abnormality of the cardiac contour;^{5, 8, 10} however, these features are not distinctive and may be absent. The key to diagnosing a coronary fistula is proper identification of an anomalous communication between the donor coronary artery and the recipient cardiac chamber or structure. Since most fistulas result in a left-to-right shunt and increased blood flow, the coronary artery will often be dilated and tortuous as in our case. Doppler color flow mapping,^{12, 14-16} transesophageal or transthoracic echocardiography,^{16, 18} contrast-enhanced electron beam tomography,^{18, 19} magnetic resonance imaging,^{16, 20} and radionuclide cardioangiograms¹³ have been proven to accurately locate the sites of origin and drainage of coronary artery fistulas. However, the presumptive gold standard is still coronary angiography.^{4, 8, 15} With the increased performance of thoracic MR, especially cardiac MR, it is important to be aware of this lesion as a potential additional source of chest pain in younger adults, typical of the active duty military population.

Surgical ligation should be considered for patients with hemodynamically significant coronary artery fistulas.^{1, 10} With age, worsening of a left-to-right shunt can result in the complications of coronary steal, congestive heart failure, angina, and endocarditis.^{2, 4, 12}

References

1. Yamanaka O, Hobbs RE. Coronary artery anomalies in 126,595 patients undergoing coronary arteriography. *Cathet Cardiovasc Diagn* 1990;21:28-40.
2. Sapin P, Frantz E, Jain A, Nichols TC, Dehmer GJ. Coronary artery fistula: an abnormality affecting all age groups. *Medicine* 1990;69:101-113.
3. Vavuranakis M, Bush CA, Boudoulas H. Coronary artery fistulas in adults: incidence, angiographic characteristics, natural history. *Cathet Cardiovasc Diagn* 1995;35:116-120.
4. Levin DC, Fellows KE, Abrams HL. Hemodynamically significant primary anomalies of the coronary arteries. Angiographic aspects. *Circulation* 1978;58:25-34.
5. Lowe JE, Oldham HN, Sabiston DC. Surgical management of congenital coronary artery fistulas. *Ann Surg* 1981; 194:373-380.
6. Friesen CH, Howlett JG, Ross DB. Traumatic coronary artery fistula management. *Ann Thorac Surg* 2000;69:1973-1982.
7. Maitre B, Jouveshomme S, Isnard R, Riquet M, Pavie A, Derenne JP. Traumatic coronary-pulmonary artery fistula, 23 years after a stab wound. *Ann Thorac Surg* 2001;70:1399-1400.
8. Wilde P, Watt I. Congenital coronary artery fistulae: six new cases with a collective review. *Clin Radiol* 1980;31:301-311.
9. Shyam Sunder KR, Balakrishnan KG, Tharakan JA, et al. Coronary artery fistulas in children and adults: a review of 25 cases with long term observations. *Int J Cardiol* 1997; 58:47-53.
10. Cheung DLC, Wing-Kuk A, Cheung HHC, Chiu CSW, Lee W. Coronary artery fistulas: long-term results of surgical correction. *Ann Thorac Surg* 2001;71:190-195.
11. Hong GJ, Lin CY, Lee CY, Loh SH, Yang HS, Liu KY, Tsai YT, Tsai CS. Congenital coronary artery fistulas: clinical considerations and surgical treatment. *ANZ J Surg* 2004;74(5):350-355.
12. Carrel T, Tkebuchava T, Jenni R, Arbenz U, Turina M. Congenital coronary artery fistulas in children and adults: diagnosis, surgical technique and results. *Cardiology* 1996;87:325-330.
13. Wells RG, Litwin SB, Sty JR. Radionuclide cardioangiographic demonstration of a coronary artery fistula. *Pediat Radiol* 1986;16:61-64.
14. Barbosa MM, Katina T, Oliveira HG, Neuenschwander FE, Oliveira EC. Doppler echocardiographic features of coronary coronary artery fistula: a report of 8 cases. *J Am Soc Echocardiogr* 1999;12:159-154.
15. Vandenbossche J, Felice H, Grivegne A, Englert M. Noninvasive imaging of left coronary arteriovenous fistula. *Chest* 1998;93:885-887.
16. Sato Y, Ishikawa K, Sakurai I, et al. Magnetic resonance imaging in diagnosis of right coronary arteriovenous fistula. *Jpn Circ J* 1997;61:1043-1046.
17. Yang Y, Bartel T, Caspari G, Eggebrecht H, Baumgart D, Erbel R. Echocardiographic detection of coronary artery fistula into the pulmonary artery. *Eur J Echocardiogr* 2001;2:292-294.
18. Brandt-Pohlmann M, Achenbach S, Pougratz G, Moshage W, Wortmann A. Non-invasive diagnosis of a congenital artery fistula. *Int J Cardiol* 1998;14:211-214.
19. Ropers D, Moshage W, Daniel WG, Jessl J, Gottwik M, Achenbach S. Visualization of coronary artery anomalies and their anatomic course by contrast-enhanced electron beam tomography and three-dimensional reconstruction. *Am J Cardiol* 2001;87:193-197.
20. Taoka Y, Nomura M, Harada M, Mitani T, Endo J, Kondo Y, Yukinaka M, Ito S, Nakaya Y, Nishitani H. Coronary-pulmonary artery fistulae depicted by multiplanar reconstruction using magnetic resonance imaging. *Jpn Circ J* 1998;62:455-457.